

# ACES: Promoting Empathy Towards Aphasia Through Language Distortion Emulation Software

Joshua Hailpern<sup>1</sup>, Marina Danilevsky<sup>1</sup>, Andrew Harris<sup>1</sup>, Karrie Karahalios<sup>1</sup>, Gary Dell<sup>2</sup>, Julie Hengst<sup>3</sup>

University of Illinois at Urbana Champaign

Urbana, IL USA 61801

<sup>1</sup>Department of Computer Science  
(jhailpe2, kkarahal)@cs.uiuc.edu  
(danilev1, harris78)@illinois.edu

<sup>2</sup>Department of Psychology  
gdell@cyrus.psych.illinois.edu

<sup>3</sup>Department of Speech and Hearing Science  
hengst@illinois.edu

## ABSTRACT

Individuals with aphasia, an acquired communication disorder, constantly struggle against a world that does not understand them. This lack of empathy and understanding negatively impacts their quality of life. While aphasic individuals may appear to have lost cognitive functioning, their impairment relates to receptive and expressive language, not to thinking processes. We introduce a novel system and model, Aphasia Characteristics Emulation Software (ACES), enabling users (e.g., caregivers, speech therapists and family) to experience, firsthand, the communication-distorting effects of aphasia. By allowing neurologically typical individuals to “walk in another’s shoes,” we aim to increase patience, awareness and understanding. ACES was grounded in the communication science and psychological literature, and informed by an initial pilot study. Results from an evaluation of 64 participants indicate that ACES provides a rich experience that increases understanding and empathy for aphasia.

## Author Keywords

Aphasia, Empathy, Emulation Software

## ACM Classification Keywords

H.5.0 General, K4.2 [Social Issues]: Assistive technologies for persons with disabilities

## General Terms

Experimentation, Human Factors

## INTRODUCTION

Receiving empathy and understanding are a constant need for those living with aphasia [17]. Aphasia is an acquired language disorder that impairs expressive and receptive language (both spoken and written)[18]. Over one million Americans are affected by aphasia [25]. There is considerable variability in the effects of aphasia, but all individuals with aphasia display difficulty producing words accurately and with ease. Specific disruptions in language include

dropping key words, substituting incorrect words for the target word, mixing up sounds or letters within words, describing items that are difficult to name, having difficulty organizing coherent strings of words, and making multiple attempts to correct such production errors.

To an outsider it may appear that an aphasic individual has poor cognitive function. However, the problem resides in the individual’s receptive and expressive language, not their ability to think. Aphasia most profoundly affects the ability to communicate with others, whose lack of understanding and empathy has the potential to “erode the social bonds that give life meaning” [17].

Our goal is to promote empathy and understanding of aphasia in unimpaired individuals. We made it possible for those without aphasia to metaphorically “walk in another’s shoes” by interacting with a system which emulates the effects of aphasia through distortion of written text. This system has the potential to help family and friends of individuals with aphasia, and to serve as a training tool for physicians, nurses, and speech-language pathologists.

We introduce a novel system and model, called Aphasia Characteristics Emulation Software (ACES), that enables users to experience the speech-distorting effects of aphasia. This model adheres to the wealth of literature in language distortions resulting from aphasia. It was informed through a series of interviews and demonstrations (using an early prototype of ACES) with professionals and students from the field of aphasia and other language disorders.

The ACES system was designed to distort a user’s Instant Messages (IMs) from the original message to one that appears like a message spoken by an individual with aphasia. Thus, the conversation that develops between the user and their IM partner has similar difficulties and hurdles to those experienced by an individual with aphasia. By experiencing these challenges firsthand, we hypothesize that users will have increased empathy, knowledge, and understanding of aphasia. Results from an evaluation of 64 participants indicate that ACES provides a rich experience that strongly increases understanding and empathy for aphasia.

The foremost contribution of our system is a demonstration of how existing literature about language distortions/errors due to cognitive impairment, age, or other language-based challenges can be extended to an interactive system that

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee.

CHI 2011, May 7–12, 2011, Vancouver, BC, Canada.

Copyright 2011 ACM 978-1-4503-0267-8/11/05...\$10.00.

increases knowledge, empathy, and awareness. Our goal is not to perfectly emulate aphasia.

## RELATED WORK

We describe aphasia, empathy and how our work builds upon, and extends, the literature related to aphasia.

### Aphasia


Aphasia is a term used to describe an acquired language disorder that is caused by damage to the left or dominant hemisphere of the brain and impairs an individual's ability to produce and understand language in both written and spoken forms [1]. The severity and pattern of aphasic symptoms vary, depending in part on the specific locations of brain damage. Clinical researchers have developed classification systems that identify different patterns or sub-types of aphasia. For example, diagnostic batteries [12,27] based on the Boston classification system are designed to categorize an individual's aphasia symptoms as either a type of non-fluent aphasia (Broca's, Transcortical Motor, Global) or fluent aphasia (Wernicke, Transcortical Sensory, Conduction, Anomic). Of particular interest to the goals of the current study is that all individuals with aphasia will display at least some difficulty with writing, and although writing may be more or less impaired than spoken language, the linguistic deficits in writing will be generally consistent with those of the person's spoken language [2]. Recent research focusing on issues of treatment and functional recovery in aphasia[5] has drawn attention to the need for clinical interventions to attend not only to the areas of deficits in the patient with aphasia, but also to the person's communicative and social systems more broadly. This paper focuses on increasing empathy for those interacting with aphasics.

### Empathy and Aphasia

Empathy is one of the fundamental underpinnings of interpersonal communication. It is an emotional response to the experiences of others, through which an empathetic person can understand and predict the feelings, emotions, and thoughts of others [10,29].

Non-technical approaches have been used to increase awareness and empathy in other situations by placing students in the "role" of an individual with an impairment. For example, students were tasked to spend a day in a wheelchair to develop an awareness of the challenges confronting a paraplegic [9]. In another course, students were given Augmentative and Alternative Communication (AAC) devices (such as text-to-speech systems which assist those with impairments or restrictions on the production or comprehension of spoken or written language) to emulate "non-vocal" communication while performing tasks such as going to a coffee shop [15].

If individuals relating to those with aphasia lack empathy and understanding, it can greatly reduce quality of life for aphasic individuals [17]. Often, family members can deny or underestimate the severity and presence of aphasic errors [6]. Further, in speech therapy, empathy is necessary to mo-



**Figure 1. Prototype Chat Window (left) and Distortion Window (right)**

*Notice how the participant's text is hard to decipher after it has been distorted by the Aphasia Model.*

*Removed (1) is a dropped Function word, Removed (2) is a random word drop and Replaced is when a word is substituted.*

tivate the aphasic client, with motivation being one of the three key aspects of effective treatment [28]. To date, research has shown that family member's ability to relate and empathize is based on how well they understand the distortions that their partners make [11].

### Technology, Empathy and Aphasia

Technology has been used as a tool to enhance the functioning and communication of individuals with aphasia [7,20,30] and other speech and language disorders [16,24]. Specifically, work in the Human Computer Interaction (HCI) community has examined technological solutions to help with day planning [23], communication [3], and increased access to web-content [8] for aphasic individuals. Most of this HCI research has focused on providing support for the individual, their communication, and providing awareness of their activities to those around them. Unlike work that focuses on developing empathic agents [21], this work aims to increase empathy, knowledge, and understanding of aphasia for those who support aphasic individuals.

### EARLY PROTOTYPE AND INITIAL INVESTIGATION

Although there is a wealth of technology that aids individuals with aphasia, to our knowledge there is none that aims to increase empathy among those associated with aphasic individuals. Following the existing work in other domains [9], we hypothesized that if individuals were to experience the effects of aphasia first hand, they might better understand the disorder and become more empathetic to its challenges.

To this end, we built a prototype chat client that randomly distorted messages sent by its user using an aphasia language model. We chose IM, because it eliminates any bias from voice inflection and focuses on communication and the language disorder itself.

To gain insight into aphasia, empathy, and the type of interaction design that may best support such a tool, we conducted an exploratory study (guided demonstrations and interviews) with ten individuals (students, professionals, and researchers) with experience in aphasia (speech and hearing science/psychology), averaging 10.3 years of experience in aphasia/language disorders, with mean age of 34.6 years.

The initial prototype (Figure 1) was built in Java and simulated an IM conversation between the user (whose text was distorted) and Eliza [31], a simple and unbiased text-based computerized conversational partner. The system provided a simplistic approximation of the effects of aphasia by randomly dropping the user's function words (16-33%), randomly dropping any words (20-40%), and randomly replacing non-function words with other random words from a dictionary file. While this prototype was a simplified representation of some effects of aphasia, it illustrated the concept and functionality of ACES in order to gather informed feedback. Users saw how their messages were distorted through an IM window, and through a log of distortions (Figure 1).

Each participant met with a researcher for one session of approximately 40 minutes duration. Sessions began with an explanation/demonstration of the prototype. Participants then used the prototype to have a conversation with Eliza. Following the IM conversation, researchers had a discussion with each participant, and concluded with a brief questionnaire using a 7-point Likert scale (7 agree, 1 disagree). There was no remuneration for participation.

### Results And Lessons Learned

Overall, participants agreed that this tool could be used to teach empathy and understanding for aphasic individuals to: friends (mean=6.1), family (mean=6.2), clinicians (mean=5.9) and professionals (mean=6.1). In addition, participants provided explicit guidance on system improvements and general suggestions on the project. For example, one participant stated that:

*This could be shown to people outside of the community of communication experts and their clients. It could teach empathy and acceptance to a group/community as a whole. -P4*

Another participant stated that this software was critical for training clinicians:

*I don't think treatment will be successful at all if the clinician is not understanding! They are the individuals who should be, and without that empathy, few gains will be achieved. -P5*

Another participant echoed P5 by stating that:

*I believe that as a simulation of a disorder... this software could be used in classrooms as early as undergrad. -P3*

All participants stated the need for a customizable future system. Given the spectrum of aphasic disorders and the manifestation variety within each sub-disorder, such a tool needs to be robust enough to emulate different types of aphasia. Participants also felt that while the emulation would not need to be 'spot on,' it would need to emulate the key distortions of each condition reasonably well.

Participants suggested applications for an aphasia emulation system. One professional speech language pathologist suggested political applications, such as helping to raise aware-

ness to increase both funding and accessibility. Another felt there to be a need for greater empathy by physicians and nurses in hospital and ER settings, who can seem dismissive of patient struggles. Most felt that an emulation system could be used by therapists with families that have a member with aphasia. All saw direct application of the software in the classroom. Some suggested that students use the software in class. Others suggested use at home, while attempting to communicate with friends for one evening. Participants mentioned that current practice in speech and hearing classrooms is to "role play" the effects of the aphasia (similar to acting). They reported that this approach is often awkward and inaccurate, indicating that an emulation system would be a large improvement over the status quo.

Perhaps the most tangible benefit of our study was the direct resources our participants provided. Students, faculty and professionals each mentioned key aspects of aphasia that should be emulated, if this system were to be useful.

### ACES: APHASIA EMULATION SOFTWARE

ACES is a both an IM client and an instantiation of a model of linguistic distortions caused by aphasia. ACES distorts messages in a manner similar to those of an individual with aphasia. Our model determines the specific nature and rate of distortions, based upon feedback from study participants and the large body of related literature. In the following sections we describe the underlying model of aphasia distortions and the user interface components of ACES. While ACES was modeled on aphasia, our system could, in principle, be used to emulate other linguistic disorders.

Please note that the contribution of our work is to promote empathy and understanding in people through experiencing the linguistic challenges of aphasic individuals. It is not to perfectly emulate aphasia. Moreover, this project focuses on expressive language distortion, and does not address those with receptive language distortion.

### Modeling The Effects of Aphasia

The effects of aphasia can vary widely based on the severity of the impairment, the type of aphasia, and even on the type of word that is subject to error. Therefore, we constructed a modeling system, Aphasia Characteristics Emulation Model Editor (**ACE-ME**), providing controls that allows the user to configure the degree and type of distortions that will be applied to their messages (Figure 2).

ACE-ME sub-divides the distortions into 5 conceptual categories: Distortion of Content Words, Distortion of Inflections, Distortion of Function Words, Distortion of Fluency, and Other. Each category defines error types that affect similar types of words (e.g., only function words or only content words). By grouping similar errors together, we leverage existing literature that focuses on specific distortion types and provide an intuitive interface for users.


ACE-ME includes the breadth of distortions experienced by individuals with aphasia. Although not every aphasic linguistic error is included, our goal was not to create a linguis-

tically exhaustive emulation tool, but one that was communicatively disruptive in a manner similar to those with aphasia. As knowledge of aphasia and natural language processing expand, so can the capabilities of our system.

### Distortion of Content Words

Most forms of aphasia affect the production of content words (verbs, nouns, adjectives, and adverbs). However, the effect and frequency of each error type differs depending upon the disorder and severity. Schwartz [26] highlights five key ways that content word production can be distorted: Formal Errors, Nonword Errors, Semantic Errors, Unrelated Errors and Mixed Errors. ACE-ME implements the first four error types. We omit Mixed Errors due to their technical complexity and low incident rate.

- Formal errors occur when a user mistakenly says a content word that is phonetically similar, yet semantically different [26]. For example: intending to say ‘population’ but stating ‘pollution.’ Formal errors are common with most aphasic individuals. ACE-ME emulates formal errors by utilizing a spell check system, JaSpell [4]. We force JaSpell to return an alternative for a word users enter (even though spelled correctly). We then randomly select one of the possible alternatives as a replacement. By using a spell check system, the word replaced would be similar to the original word given that suggested “alternatives” are based on the spelling of the original word.
- Nonword errors occur when a generated word is not a valid English word [26]. The generated nonword can be phonetically similar to the source word. For example: the word ‘castle’ is changed to ‘kaksel.’ Nonword errors are common for many aphasic speakers, particularly those with conduction or Wernike’s aphasia. ACE-ME emulates a nonword error by randomly replacing a random number of letters in the original word. Vowels are replaced by other vowels, and consonants are replaced by consonant digraphs (e.g., ‘sh,’ ‘ch’ etc) or other consonants (excluding x and z, which are uncommon in English). All generated words are then verified to not be ‘real words.’
- Semantic errors occur when the target word is replaced by another word which is semantically similar [26]. For example: ‘birthday’ is replaced with ‘anniversary,’ or ‘cake’ with ‘bread.’ While anniversary is not a synonym for birthday, nor bread for cake, they are associated. This error occurs with all aphasic types. Because semantic errors are broader than synonyms, using a thesaurus would not capture the nuance of semantic errors. Therefore, ACE-ME identifies the root of the original word [19], searches the root in a ConceptNet database [14], and chooses a random word from possible semantic matches. ConceptNet is a NLP project that, among other things, attempts to group words together based on their semantic similarity. We extracted a relational database linking words to lists of semantically related terms.
- Unrelated errors are valid English words, that have no semantic or strong phonetic relationship to the original



**Figure 2. ACES Model Editor - Distortion of Word Tab**

The overall correctness set to 61%, but the user has manually set the nonword errors to have a higher probability of occurring, increasing the overall probability of error to be 44.13%

word [26]. These errors occur particularly in severe cases of aphasia. ACE-ME randomly selects another content word from a list and replaces the original word with the unrelated term.

- Mixed errors are similar to formal errors, except the target words are semantically similar [26]. For example: ‘start’ is distorted to ‘stop,’ or ‘snail’ to ‘snake.’ Mixed errors are conceptually similar but programmatically very difficult. While we can readily generate semantically or phonetically similar words, we do not possess a large enough data set of either semantically or phonetically similar words to have a suitable intersection of the two. These errors are not very common, so not generating them should have little impact.

To understand how severity of aphasia correlates with frequency of word errors, Schwartz [26] performed an analysis of errors made in picture naming tasks. Aphasic subjects were shown a picture and asked to identify it. Schwartz examined the errors and created regression models to predict how likely a given error was to occur based on a subject’s overall level of impairment. Using the raw data published in Schwartz’s paper, we re-generated the probability of each of the five error types based on subject impairment. We used these probabilistic models in ACE-ME to calculate estimated frequency of each error. The user can move a “correctness” slider to set the probability of each of the five content word errors based on Schwartz’s work. ACE-ME also allows the user to manually set each specific content word distortion (Figure 2). ACE-ME displays the total probability of any content word error, calculated by summing the probabilities of all possible content word errors. If the user has not manually adjusted an error rate, the probability of content word error is one minus the correctness percentage.

### Distortion of Inflections

Grammatical inflection is the modification of a word to express different grammatical categories such as tense, mood,

voice, aspect, person, number, gender and case. English inflections are usually suffixes, such as the plural inflection on nouns (“apple” versus “apples”).

ACE-ME distorts only verb inflection, the most common type of inflection error for aphasic individuals (especially for those with agrammatic speech). An example of such a distortion is changing “running” to its base form, “run.” ACE-ME emulates verb inflection distortion by stemming verbs [19] and using the verb’s infinitive form. Users can set the probability of Inflection Distortion with an interface slider labeled “Inflection Morphology of Verbs.”

#### *Distortion of Function Words*

A common error made by aphasic individuals (especially those with agrammatic aphasia) is to omit function words. Function words are pronouns, articles, conjunctions, auxiliary verbs, interjections, particles, expletives, and prepositions. ACE-ME removes randomly selected function words. Users can set the probability of Function Word Distortion with an interface slider labeled “Dropping Function Words.”

#### *Distortion of Fluency*

The fluency of sentence construction can also be affected by aphasia. ACE-ME targets three common types of distortions that can affect sentence fluency: Conduit d’approche, Omissions, and Semantic Description.

- *Conduit d’approche*: This commonly affects those with conduction aphasia. Conduit d’approche occurs when an individual adds/removes or distorts the production of a word, then iteratively repairs the errors until arriving at the intended production [13]. For example: a series of Conduit d’approche would be “brepple,” “gresles,” “gla-sles,” and then completing the effort with “glasses.” ACE-ME emulates Conduit d’approche errors by repeatedly applying a random number of nonword errors to the same word. The errors are then replayed in reverse order in the sent message, giving the appearance of correcting the specific word. Each attempt is put in a separate message followed by ellipses. The slight pause gives the impression that the sender is attempting to correct the word. Once the word is “correct,” the remainder of the message is sent.
- *Omissions*: Most aphasic individuals omit or drop words from their sentences [18]. To emulate the effects of random omission, ACE-ME randomly drops words from a user’s message.
- *Semantic Description*: Many aphasic individuals can describe the features of a target word when they cannot recall the specific word [18]. This is a semantic description error (unlike a semantic error which is a one-to-one substitution). For example, an individual trying to describe glasses may say, “seeing, head, face, eyes, round.” ACE-ME utilizes the ConceptNet database [14], to search for a random number of semantically related terms and replaces the original word with a set of semantic descriptors (separated by ellipses).

Users can set the probability of all errors related to Fluency using ACE-ME interface sliders. As there is more than one type of Fluency error, ACE-ME also displays the total probability of a Fluency error occurring.

#### *Other*

There are two other effects common to aphasia that ACE-ME emulates. These are pauses and distortion awareness. Aphasic individuals often pause at atypical times. ACE-ME emulates these interruptions by randomly breaking up a message, sending each part as a separate IM.

People with certain types of aphasia (e.g., Wernicke’s) can be unaware of the errors in their speech, while others (e.g., Broca’s) are generally aware of their errors. To allow users to experience the frustration of not knowing why their conversation partner is confused (by not seeing the distorted message), as well as the frustration of not being able to send the intended message (e.g., seeing the distorted text), ACE-ME allows the user to toggle between displaying the original or the distorted message.

Users can set the probability of pauses by using the interface slider. Distortion awareness can be toggled by an interface check box in ACE-ME.

#### **ACES Interface Components**

Our system has three main components: an IM Window for engaging in instant message conversations; a Model Editor for configuring the distortions; and an Admin Window for login and switching between the IM and the Model Editor. Upon launch, users are presented with an Admin Window. Users can input a user name, password, and subject ID (used for logging all conversations to protect privacy). Users can then launch the other two components.

The ACES Model Editor (Figure 2) consists of five tabs corresponding to the five categories of distortions. Users can switch between tabs and interact with all sliders and checkboxes. All changes made to the model editor affect the current IM conversation in real-time.

The IM Window (Figure 3) is a standard IM interface with a message history, text input field, send button, and buddy list. IM history is color coded for easy identification of message origin (red=user, blue=conversation partner).

#### **Flexible Implementation**

Our system leverages a plug-in architecture for distortions. Researchers, instructors, or therapists can create or customize a specific distortion type. The ACES system examines a directory where the distortion descriptions reside, including any new distortion from this directory.

ACES sends IMs over the AOL Instant Message network, though it can easily be extended to other protocols, because it leverages the JBuddy library [32] to facilitate connection with the AOL servers. JBuddy also supports ICQ, MSN, Yahoo, Google Talk, XMPP (Jabber), Lotus Sametime, Microsoft OCS 2007, and LCS 2005.

ACES was implemented using Java 1.6, allowing the software to execute on nearly any machine, though our system has only been tested on Apple's OSX. As a result, teachers, medical staff, and researchers need not invest in new hardware to run ACES.

### ACES Logging


ACES is also conducive to post-conversation analysis and reflection because it logs all conversations in an HTML file of the perceived conversation (the user's sent messages), and in an XML file of the conversation (including the original message, the distorted message, and which distortion were applied). The HTML version is ideal for analyzing the conversation and reflecting upon difficulties in the user experience. The XML file allows researchers to analyze how users changed their behavior while trying to communicate with aphasic distortions.

### USER STUDY

In order to observe the effects of using ACES on awareness and empathy, we conducted an in-depth user study. Sixty-four individuals (grouped in subject-pairs) engaged in IM conversations with each other. We wanted to see whether experiencing text distortions first-hand through IM conversations had an impact on a subject's level of empathy for individuals with aphasia. To test this, we utilized a between-subject 2x2 factorial design. The first factor in our factorial design compared a **Treatment Group** (where subjects experienced distortions first-hand) to a **Control Group** (where subjects did not experience aphasic distortions). The second factor in our factorial design compared subjects who had prior knowledge of aphasia, through formal education or personal exposure (**Informed Sub-population**) with subjects who did not (**Uninformed Sub-population**). Our *Independent Variable* was whether or not participants experienced distortions by ACES. All participants (both treatment and control groups and both informed and uninformed sub-populations) experienced the same experimental protocol, as they were given the same prompts. Therefore subjects were blind with respect to which group (treatment or control) they were assigned. Both subjects in any given subject-pair were from the same sub-population (Informed/Uninformed) and same group (Treatment/Control).

### Experimental Protocol

Each study session consisted of four steps lasting a total of 45-60 minutes (Table 1). A session involved IMs between two participants (a subject-pair) who did not know each other. Subject-pairs remained physically separated. Each member of a pair was placed in separate identical rooms with a 23" iMac computer. All questionnaires were digital and administered using third-party software.



**Figure 3. ACES Instant Message Window**

The current (red) user's partner (blue) has their text distorted.  
 Note: this is a log from an actual experiment conversation

All participants completed both a demographic survey and a pre-study questionnaire to assess empathy using Mehrabian's measure<sup>1</sup> of emotional empathy [22]. Each participant also answered questions about their knowledge and background on aphasia. This allowed us to ensure equal background, pre-knowledge, and general empathy level across treatment and control groups. Pre-study demographic and background measures were not utilized to assess ACES effect on awareness and empathy.

Upon completion of the pre-study questionnaires, all participants were given an identical explanation of the study protocol. All participants (regardless of group or sub-population) were told that they would have two ten-minute IM conversations with each other.<sup>2</sup> During these conversations one member of the subject-pair would take on the **Aphasia Role**, while the other member of the subject-pair would take on the **Typical Role**. After the first conversation, they would switch roles (ensuring that each participant would play both roles). All subjects (in both control and treatment groups) were told that the participant in the Aphasia role would have his/her text distorted by ACES, as though they had aphasia.

	Description
<b>Step #1</b>	Demographic Questions Empathy Evaluation Pre-Study Questions on Aphasia
<b>Step #2</b>	Conversation with Partner, Prompt #1
<b>Step #3</b>	Conversation with Partner, Prompt #2 Participant Roles are Switched
<b>Step #4</b>	Final Set of Questions

**Table 1. Study Session Order**

<sup>1</sup> Mehrabian's quantitative measure of empathy assigns an individual an empathy score based on responses to 33 questions. Scores range from -132 to +132. According to Mehrabian, a representative population should yield a mean score of 33 and S.D. of 24.

<sup>2</sup> Participants were logged into IM accounts created for this study, not requiring subjects to disclose their own user names or passwords.

To guide their conversations, participants were instructed to have a “debate” or “discussion” around a specific topic. Further, each participant was assigned a position (**Pro** or **Con**), and was provided a suggested list of talking points to support their position (participants could also use their own talking points). Subjects were assigned one topic for the first conversation, and another topic for the second. The two debate topics were:

- *The age at which people gain the right to vote should be lowered to 16 years*
- *Smoking should be banned in all public places*

To conclude the study explanation, all participants were told “Remember aphasia can be mild to severe, creating distortions that are obvious, to those which are not so easily noticed or apparent.” This statement was included in the instructions so participants in the “control” group would not be concerned if they did not notice any obvious distortions.

At the end of both IM conversations, participants were administered a single questionnaire to gauge their reaction to ACES and aphasia. Participants were then remunerated with a \$10 gift certificate to Amazon.com.

### Dependent Measure

A single post-study questionnaire was utilized to assess the effects of using ACES on awareness and empathy. Ten questions on the post-study questionnaire were scored on a 7-point Likert scale. Additionally short-response questions were included to obtain qualitative reactions to ACES.

### Factorial Design, Counterbalancing, Treatment Effects

Our cohort of sixty-four individuals consisted of two sub-populations. Thirty-two subjects (one half of our subjects) had prior knowledge of aphasia either through formal education or personal exposure (Informed Sub-population), and the other thirty-two subjects did not (Uninformed Sub-population). Further, half of each sub-population was in the Control Group, and half was in the Treatment Group. Thus, we used a **2x2 Between-Subject Factorial Design** with 16 subjects (8 subject-pairs) in each of the following **Experimental Conditions**: Informed-Treatment, Informed-Control, Uninformed-Treatment and Uninformed-Control. By testing both a Informed and Uninformed population, we had the ability to test the potential applicability of ACES to both family members and clinicians/doctors in training.

Given the number of other factors inherently present in this type of experimental setup, there were at least three potential confounding effects (debate topic order, debate position and role order). To help control for these potential confounding effects, we used *counterbalancing*, a method commonly employed to help avoid confounding from order of task and presentation. When counterbalancing, all permutations of the confounders are included in an attempt to minimize any bias due to these confounders that are not central to the experimental question. Thus, for each Experimental Condition (e.g., Informed/Uninformed and Treatment/Control), we

	<b>% Male</b>	<b>Age (SD)</b>	<b>Empathy<sup>1</sup> (SD)</b>
<b>Overall</b>	39.06	26.05 (7.30)	34.52 (25.82)
<b>Overall - C</b>	28.12	25.06 (7.27)	40.09 (22.60)
<b>Overall - T</b>	50.00	27.03 (7.30)	28.94 (27.92)
<b>Informed</b>	37.50	25.35 (6.83)	39.25 (25.38)
<b>Informed - C</b>	25.00	24.13 (6.54)	46.81 (21.28)
<b>Informed - T</b>	50.00	26.56 (7.11)	31.69 (27.51)
<b>Uninformed</b>	40.62	26.75 (7.78)	29.78 (25.76)
<b>Uninformed - C</b>	31.25	26.00 (8.04)	33.38 (22.50)
<b>Uninformed - T</b>	50.00	27.50 (7.69)	26.19 (28.95)
	<b>Highest Level of Education Completed (%)</b>		
	<i>High School</i>	<i>Bachelors</i>	<i>Graduate</i>
<b>Overall</b>	25.00	56.25	18.75
<b>Overall - C</b>	28.12	56.25	15.63
<b>Overall - T</b>	21.88	56.25	21.87
<b>Informed</b>	34.38	53.12	12.50
<b>Informed - C</b>	43.75	37.50	18.75
<b>Informed - T</b>	25.00	68.75	6.25
<b>Uninformed</b>	15.62	59.38	25.00
<b>Uninformed - C</b>	12.50	75.00	12.50
<b>Uninformed - T</b>	18.75	43.75	37.50

**Table 2. Population Demographics & Empathy Scores**

*Empathy refers to Mehrabian’s measure of emotional empathy<sup>3</sup>.*

*Mean age and empathy scores are shown.*

*C = Control T = Treatment*

needed 8 pairs of conversations to account for all permutations of Role (Aphasic/Typical), Topic (Voting/Smoking), and Debate Position (Pro/Con).

In addition, we attempted to control for “treatment effects.” A treatment effect occurs when a participant has a reaction to a placebo simply because they are told they are going to receive a treatment. By effectively telling both Treatment and Control groups that they were going to be in the “treatment group,” we attempted to control for the treatment effect. If we noticed a positive or negative response from participants in our control group, we would have confidence that it would be due to treatment effects, thus allowing us to better qualify the effects reported in the treatment group. Likewise, if no effects were observed in the control group, we would feel confident that the simple act of using IM in this experiment did not contribute to those results, and were due directly to ACES emulating the effects of aphasia.

### Statistical Tests

To examine the effects of ACES we compared responses from the four Experimental Conditions to the post-study questionnaire in Step 4 (Table 1). A 2-way ANOVA (analysis of variance) was used to test for differences between the four Experimental Conditions (Group by Sub-Population). An ANOVA test assumes that data are normally distributed and is robust to deviations in normality. Data is presented as median, inter-quartile range (the range of values that span the middle 50% of the data).

	Control Group		Treatment Group		2-Way ANOVA	
	Median	Mean (SD)	Median	Mean (SD)	F Statistic	p -val
1 In hindsight, this experience made me <> empathetic to individuals with aphasia.†	5 [4,5]	4.75 (0.80)	6 [6,6]	5.91 (0.77)	F(1,61)=34.84	0.0001
2 In hindsight, this experience made me <> empathetic to people who try to understand an individual with aphasia. †	4 [4,5]	4.62 (0.87)	6 [5,6]	5.59 (0.87)	F(1,61)=19.72	0.0001
3 This tool is useful for building empathy for family members‡	4 [4,5]	4.19 (1.47)	6.5* [5,7]	6.09 (1.06)	F(1,61)=34.96	0.0001
4 This tool is useful for building empathy for speech therapists‡	4 [4,5]	4.47 (1.59)	6 [5,7]	6.09 (1.03)	F(1,61)=23.28	0.0001
5 This tool is useful for building empathy for nurses and doctors‡	4.5* [4,5]	4.47 (1.54)	7 [5,7]	6.22 (0.91)	F(1,61)=30.13	0.0001
6 This tool is useful for building empathy for care givers‡	4 [4,5]	4.38 (1.50)	7 [6,7]	6.38 (0.91)	F(1,61)=41.12	0.0001
7 This tool is useful for education/training of individuals who treat/interact with individuals with aphasia‡	4 [4,6]	4.5 (1.65)	7 [6,7]	6.53 (0.80)	F(1,61)=39.02	0.0001
8 Which role helped you understand aphasia better?§	4 [3.5,4]	3.91 (1.42)	2 [1,3]	2.28 (1.55)	F(1,61)=18.79	0.0001
9 Which role increased your empathy for the the perspective of understanding an individual with aphasia?§	4 [4,4.5]*	4.5 (1.03)	1.5* [3.5,6]*	3.72 (2.40)	F(1,61)=1.02	0.317
10 Which role increased your empathy for the the perspective of an individual with aphasia?§	4 [3.5,4]*	3.69 (1.38)	2 [1,4.5]	2.72 (2.25)	F(1,61)=4.26	0.043

**Table 3. Results**

*Median represents median value and interquartile range [IQ range] - 2-Way ANOVA Values represents Treatment vs. Control Factor*

*2-Way ANOVA Values for Informed Sub-population vs. Uninformed Sub-population (not shown) all  $p > 0.05$*

\* Half values are due to even number of data points where the value separating the higher half from the lower half lies between two different values, resulting in a median which is the average of the two values. For example a data set of [1,4,4,5,5,7] would have a median of 4.5 although 4.5 is not a possible value.

† Question Choices Were: 1=Much Less, 2=Less, 3=Slightly Less 4= No Change, 5= Slightly More, 6=More, 7=Much More

‡ Question Choices Were: 1= Disagree, 4=Neutral, 7 = Agree § Question Choices Were: 1= Aphasia, 4=Equal, 7=No Distortions/Normal

## Subjects

Sixty-four participants completed the ACES experiment (Table 2). More than 75% of the Informed population had an educational background in psychology, speech and hearing science, or traditional sciences. Participants in the Uninformed population came from a wide variety of areas spanning engineering, science, and liberal arts.

The empathy scores of our subject population, as measured at the beginning of each session (Table 2), closely follow those of a standard population as presented in the original paper by Mehrabian[22]. With a balanced male/female population, Mehrabian expects a mean score of 33 (standard deviation of 24). Our study cohort had a mean empathy score of 34.5 (standard deviation of 25.8). This is nearly identical to Mehrabian. To ensure control and treatment groups were equivalent, we examined the empathy scores of the groups and the four initial self-response questions regarding participant knowledge of and sensitivity to the challenges of communication with aphasia. We examined any differences overall and within the Informed/Uninformed demographics. There were no statistically significant differences ( $p = 0.08$ ,  $z=1.75$ ) between the control and treatment groups in a priori disposition, knowledge of, or empathy towards individuals with aphasia (results not shown).

## Results

Overall, participants in the group experiencing aphasic distortions by the ACES were strongly affected by their experience relative to the control group (Table 3, Questions 1 & 2). For example, participants indicated that the experience made them more empathetic to individuals with aphasia compared to the control group who felt only a slight change, with  $p < 0.0001$ . Participants in the treatment group strongly

agreed that ACES should be used in all applications explicitly listed, while the control group felt neutral as to the applicability of ACES (Table 3, Questions 3-7). Participants in the Treatment group strongly agreed that ACES could be used to increase empathy in caregivers, while participants that did not experience distortions felt neutral towards ACES use, with  $p < 0.0001$ . Participants in the Treatment group felt that they gained understanding and empathy from having their text distorted, while the control group did not rank one role higher than another (Table 3, Questions 8 & 10).

When data was analyzed separately for the Informed and Uninformed groups, results were similar to those of the full cohort (results not shown). Finally, we compared the Informed treatment group with the Uninformed treatment group to see if a priori knowledge of aphasia influenced subject response. Both treatment groups found ACES beneficial in increasing knowledge and empathy.

Though not shown in Table 3, there was no significant difference in any of the post-study questions by Informed/Uninformed Sub-Population. Furthermore, there were no interaction effects between the Treatment Group (Treatment/Control) and Sub-Population (Informed/Uninformed) (data not shown - available from the authors upon request).

Figure 3 is a representative conversation between two subjects (taken from our study). The IM window is from jet587, whose partner, apple36, is having his text distorted.

## Post Hoc Results

The one question with a non-significant value overall was question 9: "Which role increased your empathy for the the perspective of understanding an individual with aphasia?" Given the large inter-quartile range within the Treatment

group, we hypothesized that the lack of significance was due to learning effects (having more exposure to the intervention makes you increasingly more empathetic). To test this hypothesis, we subdivided the Treatment Group by comparing the responses of subjects who were in the Aphasic Role first, to those in the Typical Role first. Because these are mutually exclusive subsets of the Treatment group we performed a between subject analysis comparing responses with a unpaired student T-test and found  $p=0.015$  ( $-2.66=t$ ). In particular, the mean response ( $2.69 \pm 1.99$ ) of participants whose first role was Typical, indicated that the Aphasic role increased empathy more. In contrast, the mean response ( $4.75 \pm 2.38$ ) of participants whose first role was Aphasic, indicated that the Typical role increased empathy more.

Thus, whichever role the participant experienced second was the role reported to have increased their empathy more. This post-hoc analysis suggests that it does not matter which role (Aphasic or Typical) participants take on first. Rather, we hypothesize that it is the opportunity to play both roles that may increase empathy. This would need to be validated in future experiments. We tested the other questions for learning effects and did not find any.

## DISCUSSION

Participants who experienced the distortions of aphasia had a much stronger response than those in the control group. Those participants in the treatment group reported “strong” effects from the experiment, including increased empathy and envisioning a wide variety of potential uses of ACES. In contrast, participants who did not have their text distorted reported little or no change in their perception of or empathy towards aphasia. The median quantitative responses from the control group were almost unanimously at 4 (a neutral opinion/no change), with little deviation in inter-quartile ranges. The control group also saw no application for ACES to increase empathy or awareness of aphasia. This is in stark contrast to those with exposure to aphasic distortions.

There is always the possibility of a change in empathy resulting from being in an experimental condition (known as experimental demand). However, given the great difference in responses between the two groups (qualitatively and statistically), we are confident that ACES demonstrates a real impact on increasing awareness and empathy by experiencing distortions caused by aphasia. This is further supported in that both groups were given identical prompts, and the experiment was fully counterbalanced.

While quantitative feedback provides strong evidence of the effectiveness of ACES, qualitative feedback highlights how informative the ACES experience was for participants, and how supportive they are of the project. For example, one participant from the Informed Treatment group stated:

*This was a wonderful way to gain empathy! I've learned about aphasia in classes, but this perspective was very helpful and will make a lasting impression.*

The qualitative feedback supported how prior experience with aphasia was not a prerequisite for gaining insight. One participant from the Uninformed Treatment group stated:

*It was the most eye-opening from the point of view of having my text distorted. It was amazing how hard it can be sometimes to get a point across. From the aphasic side it is almost like experiencing it from both sides, because you see how difficult what you say can be for the 'normal' person.*

Participants also suggested that even short exposure could make a lasting impact, saying, “it would only take a few minutes to gain perspective.”

In contrast, qualitative feedback from subjects in the control group supported our quantitative findings, indicating that this was not a meaningful experience. Given the stark difference in qualitative and quantitative responses between the control and treatment groups, we believe that ACES can provide a meaningful experience for all users who seek to better understand and empathize with aphasic individuals.

## FUTURE WORK

Our experiment examined the impact of ACES on participants' empathy and understanding of aphasic individuals. However, our long term goal is to meaningfully improve real-world interactions, thereby improving quality of life and therapy for individuals with aphasia. We propose conducting a follow-up study to explore whether a subject's experience with ACES can positively impact their real-world interactions with individuals with aphasia. This study would directly involve individuals with aphasia in order to determine if they can perceive an increase in empathy present in users of ACES. We further wish to investigate the long-term impact of ACES, and how to integrate the software within a classroom of therapeutic setting

While ACES supports a robust set of distortions, there are always refinements and less common errors that can be implemented. Over time, we hope to increase the capabilities and functionality that ACES provides.

Though this project explicitly targets aphasia, our goal is to introduce language distortion emulation as a new approach to increasing empathy for those with other language impairments. We envision, for instance, a system similar to ACES that distorts text as though it comes from a young child. This can teach patience, understanding, and empathy to parents, teachers, and caregivers.

## CONCLUSION

Empathy and understanding from family members, friends and clinicians can enhance the quality of life of aphasic individuals. Family members can deny or underestimate the severity and presence of aphasic errors. Without empathy, the quality of speech therapy can suffer, jeopardizing the speed and recovery of aphasic individuals. Our work has made several contributions to address these concerns stemming from a lack of empathy.

First, we leveraged speech-language and psychological theory to design and construct a model of aphasic distortions. Second, from an initial investigation we refined our model and designed a system to increase awareness and empathy with aphasic individuals. Third, we developed a novel system (ACES) that allows a neurologically typical individual to experience firsthand, the linguistic distortions of aphasia. Fourth, ACES was validated in an investigation with 64 participants (with and without background on aphasia). Results from this study strongly show that using ACES can increase empathy and awareness of aphasia.

Our model and system demonstrate how technology can play a central role in increasing empathy, awareness and understanding for individuals with a language disorder. Our approach can be used in many other domains where atypical language is present and can be emulated. It is through empathy that we learn to understand each other.

#### ACKNOWLEDGMENTS

We would like to thank our reviewers, participants, family and friends.

#### REFERENCES

- Benson, D. F. *Aphasia, Alexia and Agraphia: Clinical Neurology and Neurosurgery Monographs*. Churchill Livingstone, New York, 1979.
- Benson, D. F. *The neurology of thinking*. Oxford University Press, USA, 1994.
- Boyd-Graber, J. L. et al. Participatory design with proxies: developing a desktop-PDA system to support people with aphasia. In *Proceedings of the CHI 06 (2006)*. ACM, Montreal, Canada.
- Bruno Martins. JaSpell. <http://jaspell.sourceforge.net/>.
- Chapey, R. *Language intervention strategies in aphasia and related neurogenic communication disorders*. Lippincott Williams & Wilkins, 2001.
- Czvik, P. S. Assessment of family attitudes toward aphasic patients with severe auditory processing disorders. In *Proceedings of Clinical Aphasiology (1977)*.
- Daemen, E. et al. Designing a free style, indirect, and interactive storytelling application for people with aphasia. *INTERACT 2007*, 2010, 221-234.
- Devlin, S. and Unthank, G. Helping aphasic people process online information. In *Proc. of Assets '06 (2006)*. ACM, Portland, Oregon, USA New York, NY, USA.
- Doerksen, S. *Recreation for Persons with Disabilities (RPM 277)*. Pennsylvania State University Department of Recreation, Park and Tourism Management, Fall, 2009.
- Dymond, R. F. A scale for the measurement of empathic ability. *Journal of Consulting Psychology*, 13(2), 1949, 127-133.
- Furbacher, E. A. and Wertz, R. T. *Simulation of Aphasia by Wives of Aphasic Patients*. *Clinical aphasiology*, 1983, 227.
- Goodglass, H., Goodglass, Kaplan. *Boston Diagnostic Aphasia Examination: Stimulus Cards--Short Form*. Lippincott Williams & Wilkins, 2001.
- H. Goodglass and Kaplan, E. *The Assessment of Aphasia and Other Neurological Disorders*. Williams and Wilkins, Baltimore, MD, 1983.
- Hugo Liu, Aingh, Push, and Eslick, Ian. ConceptNet. <http://web.media.mit.edu/~hugo/conceptnet/>.
- Hengst, J. *Augmentative and Alternative Communication (SHS 473)*. University of Illinois Department of Speech and Hearing Science, Spring, 2005.
- Kitzing, P., Maier, A., and Åhlander, V. L. Automatic speech recognition (ASR) and its use as a tool for assessment or therapy of voice, speech, and language disorders. *Logopedics Phoniatrics Vocology*, 34(2), 2009, 91-96.
- Liechty, J. A. and Heinzekehr, J. B. Caring for Those Without Words: A Perspective on Aphasia. *The Journal of Neuroscience Nursing*, 39(5), 2007, 316.
- Sarno, M. T. *Acquired aphasia (Third Edition)*. Academic Press, San Diego, CA, 1998.
- Martin Porter. Snowball: A language for stemming algorithms. 2010, <http://snowball.tartarus.org/>.
- McCall, D. et al. Integrating technology and targeted treatment to improve narrative production in aphasia: A case study. *Aphasiology*, 23(4), 2009, 438-461.
- McQuiggan, S. W. and Lester, J. C. Learning empathy: A data-driven framework for modeling empathetic companion agents. In *Proceedings of Autonomous Agents and Multiagent Systems (2006)*. ACM, New York, NY.
- Mehrabian, A. and Epstein, N. A measure of emotional empathy. *Journal of personality*, 40(4), 1972, 525-543.
- Moffatt, K. et al. The participatory design of a sound and image enhanced daily planner for people with aphasia. In *Proceedings of CHI 04 (2004)*. ACM, NY.
- Mullennix, J. and Stern, S. *Computer Synthesized Speech Technologies: Tools for Aiding Impairment*. Medical Information Science Reference, Hershey, PA, 2010.
- National Institute on Deafness and Other Communication Disorders. Aphasia. 2010, <http://www.nidcd.nih.gov/health/voice/aphasia.htm>.
- Schwartz, M. F. et al. A case-series test of the interactive two-step model of lexical access: Evidence from picture naming. *Journal of Memory and Language*, 54(2), 2006, 228-264.
- Shewan, C. M. and Kertesz, A. Reliability and validity characteristics of the Western Aphasia Battery (WAB). *Journal of Speech and Hearing Disorders*, 45(3), 1980, 308.
- Shill, M. A. Motivational factors in aphasia therapy: Research suggestions. *Journal of Communication Disorders*, 12(6), 1979, 503-517.
- Stotland, E. Exploratory investigations of empathy. *Advances in experimental social psychology*, 4, 1969, 271-314.
- van de Sandt-Koenderman, M., Wieggers, J., and Hardy, P. A computerised communication aid for people with aphasia. *Disability & Rehabilitation*, 27(9), 2005, 529-533.
- Weizenbaum, J. Contextual understanding by computers. *Communications of the ACM*, 10(8), 1967, 480.
- Zion Software. JBuddy Messenger. <http://www.zionsoftware.com/products/messenger/>.